

IN SITU SIMULATION OF IN-HOSPITAL CARDIAC ARREST TO IMPROVE RESIDENT PHYSICIAN CONFIDENCE AND PERFORMANCE

Christine Sailer, MD^a, Lynne Rosenberg, MD^b, Alexis Tumolo, MD^a, Bridget Kovach, MAEdHD^c, Christopher P. Kovach, MD, MSc^a.

^aDivision of Cardiology, Department of Medicine, University of Colorado, Aurora, CO.

^bDepartment of Medicine, University of Colorado, Aurora, CO.

^cDepartment of Professional Development, Children's Hospital Colorado, Aurora, CO.

Background: In-hospital cardiac arrest (IHCA) resuscitations at academic institutions are frequently led by resident physicians with one year of post-graduate training who have not received formal teamwork, communication, or leadership training.¹⁻³ Resident physicians who respond to an IHCA are thus called upon to implement infrequently used, non-technical skills in an emergent situation with high mental workload and an ad hoc team. Recent studies have suggested that reinforcement of ACLS skills every two years is insufficient to maintain high-quality resuscitation practice.^{1,4-6} Brief, repeated, in-situ (i.e., workplace-based) resuscitation training, however, may lead to improved skill retention.^{7, 10-14} Cardiac arrest simulation is a well-established educational strategy for enhancing learner performance in resuscitation.⁷⁻⁹

Methods: A needs assessment performed in May 2021 demonstrated that 46% of all internal medicine residents and 95% of rising PYG 2s feel unprepared to lead a cardiac arrest resuscitation. A one-hour, monthly educational intervention utilizing in-situ, scenario-based medical simulation was developed and implemented to enhance the technical, leadership, and communication skills of resident physicians providing care in the cardiac intensive care unit (Fig 1). A Prestan Series 2000 CPR Manikin and the Simpl app were purchased as a low-cost alternative to high-tech simulation manikins. Pre- and post-intervention self-assessments of resident confidence were obtained using a 5-point Likert scale questionnaire. Skill performance was assessed during the first and last simulation using a standardized rubric (Fig 2). Changes in resident confidence and performance were described and compared using a paired t-test.

Results: A total of 7 educational interventions involving 42 internal medicine residents (20 PYG 1, 21 PYG2, 1 PYG3) have been completed to date. Preliminary data from the skills assessment demonstrated significant improvement in time to first defibrillation (114 ± 44 vs 50 ± 33 seconds, $p=0.005$, Fig 3), interruptions in CPR (18 ± 5 vs 7 ± 2 % of time off chest, $p=0.001$, Fig 4) and leadership and communication score (3 of 6 vs 6 of 6, $p=0.0002$). Following the intervention, more residents reported feeling "somewhat or very confident" to lead a cardiac arrest resuscitation (31 vs 72%), provide feedback on CPR quality (41 vs 69%), order appropriate medication and specify dose (38 vs 62%), provide high-quality post-resuscitative care (34 vs 76%) and discontinue cardiac arrest resuscitation for futility (14 vs 69%) (Fig 5). 100% of respondents reported the simulation curriculum improved their understanding of cardiac arrest resuscitation (76% "strongly agree", 24% "somewhat agree").

Conclusion: An in-situ simulation curriculum was effective in improving resident physician performance in leading an IHCA resuscitation. Further studies are needed to evaluate retention of performance gains over time.

APPENDIX

Figure 1. Overview of Educational Intervention




Figure 2. Pre and Post Intervention Skills Assessment Rubric

Simulation Skills Checklist
<p>ACLS Skills</p> <p>Time to first shock: ___ seconds</p> <p>Interruptions in CPR: time off the chest ___seconds / total CPR ___seconds = ___ % time off chest</p> <p>Yes No</p> <p><input type="checkbox"/> <input type="checkbox"/> Rate 100-120 >90% of the time</p> <p><input type="checkbox"/> <input type="checkbox"/> Depth at least 2in >90 % of the time</p> <p><input type="checkbox"/> <input type="checkbox"/> Appropriate dose/administration of epi and amio</p> <p><input type="checkbox"/> <input type="checkbox"/> ECG obtained post ROSC, cath lab activated if STEMI</p> <p><input type="checkbox"/> <input type="checkbox"/> Mental status assessed, TTM initiated</p> <p>Leadership & Communication Skills</p> <p>Yes No</p> <p><input type="checkbox"/> <input type="checkbox"/> Leader identifies themselves, stands at foot of bed</p> <p><input type="checkbox"/> <input type="checkbox"/> Leader identifies team members</p> <p><input type="checkbox"/> <input type="checkbox"/> Leader concisely assigns tasks to specific individuals >90% of the time</p> <p><input type="checkbox"/> <input type="checkbox"/> Team utilizes closed loop communication > 90% of the time</p> <p><input type="checkbox"/> <input type="checkbox"/> Leader verbalizes rhythm at every pulse check</p> <p><input type="checkbox"/> <input type="checkbox"/> Leader summarizes resuscitation course periodically (i.e., every 2-3 cycles)</p>

Figure 3. Time to First Shock




Figure 4. Interruptions in CPR





Figure 5. Resident Confidence Pre- and Post-Intervention



REFERENCES

1. Merchant RM, Topjian AA, Panchal AR, Cheng A, Aziz K, Berg KM, et al. Part 1: Executive Summary: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2020;142(16_suppl_2):S337-s57.
2. Burden AR, Pukenas EW, Deal ER, Coursin DB, Dodson GM, Staman GW, et al. Using Simulation Education With Deliberate Practice to Teach Leadership and Resource Management Skills to Senior Resident Code Leaders. *J Grad Med Educ*. 2014;6(3):463-9.
3. Rosen MA, DiazGranados D, Dietz AS, Benishek LE, Thompson D, Pronovost PJ, et al. Teamwork in healthcare: Key discoveries enabling safer, high-quality care. *Am Psychol*. 2018;73(4):433-50.
4. Committee on the Treatment of Cardiac Arrest: Current S, Future D, Board on Health Sciences P, Institute of M. The National Academies Collection: Reports funded by National Institutes of Health. In: Graham R, McCoy MA, Schultz AM, editors. *Strategies to Improve Cardiac Arrest Survival: A Time to Act*. Washington (DC): National Academies Press (US). Copyright 2015 by the National Academy of Sciences. All rights reserved.; 2015.
5. Nelson J, Abella BS. Improving CPR measurement: are we there yet? *Resuscitation*. 2015;87:A1-2.
6. Yeung J, Djarv T, Hsieh MJ, Sawyer T, Lockey A, Finn J, et al. Spaced learning versus massed learning in resuscitation - A systematic review. *Resuscitation*. 2020;156:61-71.
7. Greif R, Lockey A, Breckwoldt J, Carmona F, Conaghan P, Kuzovlev A, et al. European Resuscitation Council Guidelines 2021: Education for resuscitation. *Resuscitation*. 2021;161:388-407.
8. Mundell WC, Kennedy CC, Szostek JH, Cook DA. Simulation technology for resuscitation training: a systematic review and meta-analysis. *Resuscitation*. 2013;84(9):1174-83.
9. Greif R, Bhanji F, Bigham BL, Bray J, Breckwoldt J, Cheng A, et al. Education, Implementation, and Teams: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2020;142(16_suppl_1):S222-s83.
10. Cheng A, Eppich W, Grant V, Sherbino J, Zendejas B, Cook DA. Debriefing for technology-enhanced simulation: a systematic review and meta-analysis. *Med Educ*. 2014;48(7):657-66.
11. Sutton RM, Niles D, Meaney PA, Aplenc R, French B, Abella BS, et al. Low-dose, high-frequency CPR training improves skill retention of in-hospital pediatric providers. *Pediatrics*. 2011;128(1):e145-51.
12. Sutton RM, Niles D, Meaney PA, Aplenc R, French B, Abella BS, et al. "Booster" training: evaluation of instructor-led bedside cardiopulmonary resuscitation skill training and automated corrective feedback to improve cardiopulmonary resuscitation compliance of Pediatric Basic Life Support providers during simulated cardiac arrest. *Pediatr Crit Care Med*. 2011;12(3):e116-21.
13. Oermann MH, Kardong-Edgren SE, Odom-Maryon T. Effects of monthly practice on nursing students' CPR psychomotor skill performance. *Resuscitation*. 2011;82(4):447-53.
14. Cheng A, Lockey A, Bhanji F, Lin Y, Hunt EA, Lang E. The use of high-fidelity manikins for advanced life support training--A systematic review and meta-analysis. *Resuscitation*. 2015;93:142-9.