



Children's Hospital Colorado
Musculoskeletal Research Center

Successful Conservative Management of Kienbock's Disease in a 7-year-old: A Case Report

Ryan T. Calkins, B.S.¹; James Lendrum, MD/MPH¹; Alex Lencioni, MD¹; Tyler Freeman, MD¹; Frank Scott, MD^{1,2}

¹Department of Orthopaedic Surgery, University of Colorado School of Medicine, Aurora, CO

²Department of Orthopaedic Surgery, Children's Hospital Colorado, Aurora, CO



ORTHO
University of Colorado

Abstract

Background: Kienbock's disease is primarily seen in young adult males but has been recognized in skeletally immature populations as well. Traditional treatment strategies recommend operative treatment but high remodeling potential in children may allow conservative management.

Case Description : We present the case of a 7-year-old female with two months of atraumatic right wrist pain who was found to have edematous signal change within the lunate on wrist MRI consistent with Kienbock's disease. She was treated with rigid immobilization for 12 weeks and transitioned to custom orthotic splint for another 3 months during activities. At her 6-month follow-up, she reported minimal wrist pain with repeat MRI demonstrating resolution of lunate edema.

Literature Review: Available literature shows a significant portion of patients treated conservatively subsequently require surgical intervention due to unresolved symptoms or progressive disease. Only three cases are reported in the literature where skeletally immature patients were successfully treated with conservative management alone

Clinical Relevance: We report the youngest case of Lichtman stage I Kienbock's disease successfully treated with conservative management resulting in clinical and imaging resolution. Younger patients may be able to successfully remodel and recover from Kienbock disease with extended time in conservative management.

Introduction


Kienbock's disease is primarily seen in young adult males It is rare in children and often referred to as "Teenbock's Disease" as the treatments are not as clear as with adults

Methods

7-year-old female presented with Kienbock's disease and subsequently followed-up at 8 weeks, 6 months, and 3 years post injury.

Institutional Review Board (IRB) exemption was obtained, and patient/family was informed and consented to the case report.

Case Timeline



Discussion/Conclusions

This is the youngest case of Lichtman stage I Kienbock's disease successfully treated with conservative management resulting in clinical and imaging resolution.

Younger patients may be able to successfully remodel and recover from Kienbock's disease with extended time in conservative management.

Duration of conservative treatment in this case was consistent with the recently described protocols of Lichtman et al.

This case report has limitations as it is unable to be generalized and may not apply to similar cases due to inability to understand every extraneous factor affecting disease progression and resolution in this patient.

Acknowledgements/Additional Info

Thank you to Andy Lalka, MPH for his management of this Case Report.

No funding was received for this work.

No conflicts of interest to report for any authors.

References

- van Leeuwen, W.F., et al., *Risk Factors of Lunate Collapse in Kienbock Disease*. J Hand Surg Am, 2017. **42**(11): p. 883-888 e1.
- Camus, E.J. and L. Van Overstraeten, *Kienbock's disease in 2021*. Orthop Traumatol Surg Res, 2021: p. 103161.
- Camus, E.J., L. Van Overstraeten, and F. Schuind, *Lunate biomechanics: application to Kienbock's disease and its treatment*. Hand Surg Rehabil, 2021. **40**(2): p. 117-125.
- Alshar, A., *Kienbock Disease in the Skeletally Immature Patient*. J Hand Surg Am, 2018. **43**(5): p. 465-469.
- Ando, Y., et al., *Temporary scaphotrapezoidal joint fixation for adolescent Kienbock's disease*. J Hand Surg Am, 2009. **34**(1): p. 14-9.
- Docquier, P.L., A. Traore, and S. Hugon, *Kienbock's disease in a 6-year-old boy*. Acta Orthop Belg, 2009. **75**(6): p. 832-5.
- Edelson, G., N.D. Reis, and D. Fuchs, *Recurrence of Kienbock disease in a twelve-year-old after radial shortening. Report of a case*. J Bone Joint Surg Am, 1988. **70**(8): p. 1243-5.
- Foster, R.J., *Kienbock's disease in an 8-year-old girl: a case report*. J Hand Surg Am, 1996. **21**(4): p. 595-8.
- Herdem, M., C. Ozkan, and H. Bayram, *Overgrowth after radial shortening for Kienbock's disease in a teenager: case report*. J Hand Surg Am, 2006. **31**(8): p. 1322-5.
- Hong, I.T., et al., *Kienbock's disease with non-negative ulnar variance : Treatment with combined radial wedge and shortening osteotomy*. Orthopade, 2019. **48**(1): p. 96-101.
- Hosking, O.R., *Kienbock's disease in an 8 year old boy*. Aust N Z J Surg, 1989. **59**(1): p. 92-3.
- Kazuki, K., et al., *Time course of magnetic resonance images in an adolescent patient with Kienbock's disease treated by temporary scaphotrapezoidal joint fixation: a case report*. J Hand Surg Am, 2006. **31**(1): p. 63-7.
- Mathieu, L. and C. Dumontier, *[Kienbock's disease in a nine-year-old boy: a case report]*. Chir Main, 2009. **28**(2): p. 99-102.
- Matsuhashi, T., et al., *Radial overgrowth after radial shortening osteotomies for skeletally immature patients with Kienbock's disease*. J Hand Surg Am, 2009. **34**(7): p. 1242-7.
- Rhee, S.K., et al., *A comparative study of the surgical procedures to treat advanced Kienbock's disease*. J Korean Med Sci, 1996. **11**(2): p. 171-8.
- Yasuda, M., et al., *Temporary scapho-trapezoidal joint fixation for Kienbock's disease in a 12-year-old girl: a case report*. J Hand Surg Am, 1998. **23**(3): p. 411-4.
- Shigematsu, K., et al., *Treatment of Kienbock's disease with cultured stem cell-seeded hybrid tendon roll interposition arthroplasty: experimental study*. J Orthop Sci, 2006. **11**(2): p. 198-203.
- Luo, J. and E. Diao, *Kienbock's disease: an approach to treatment*. Hand Clin, 2006. **22**(4): p. 465-73; abstract vi.
- Nealey, E.M., et al., *Radiologic Guide to Surgical Treatment of Kienbock's Disease*. Curr Probl Diagn Radiol, 2018. **47**(2): p. 103-109.
- Schuind, F., S. Eslami, and P. Ledoux, *Kienbock's disease*. J Bone Joint Surg Br, 2008. **90**(2): p. 133-9.

Initial Presentation



Figure 1: AP, Lateral and Oblique plain radiographic projections of the right wrist upon initial presentation without any signs of carpal mal-alignment or osseous change to the lunate.

Figure 2: Axial, Sagittal and Coronal T1 FS post contrast sequences of the right wrist showing edema of the Lunate.

8-Week Follow-Up



Figure 3: Anterior-Posterior, Lateral and Oblique plain radiographic projections of the right wrist after eight weeks of immobilization, without any signs of carpal mal-alignment or osseous change to the lunate.

6-Month Follow-Up



Figure 4 Axial, Sagittal and Coronal PD FS sequences of the right wrist showing resolution of the edema within the lunate at six months after initial presentation.

3-Year Follow-Up



Figure 5: Radiographs without any signs of carpal mal-alignment or osseous change to the lunate