



University of Colorado  
Anschutz Medical Campus

Mohamoud Ahmed, Yaron Firezen, Tim Vigers, Kathryn Akong, Julie Ryu, Andrea Hahn, Hani Fanous, Anastassios Koumbourlis, Pornchai Tirakitsoontorn, Antonio Arrieta, Elizabeth Burgener, Jonathan Cogen, Drake Bouzek, Elin Towler, Allison Keck, Erika Peterson, Dayna Stout, John Bradley, Scott Sagel

Children's Hospital Colorado, Rady Children's Hospital-San Diego, Children's National Hospital, Children's Hospital of Orange County, Seattle Children's Hospital



Children's Hospital Colorado

## BACKGROUND

Availability of CFTR modulator therapy down to younger ages coupled with the COVID-19 pandemic led to a remarkable decrease in pulmonary exacerbations (PEX) requiring hospitalization among children with CF

## STUDY OBJECTIVE


Determine the characteristics of children with CF who are now requiring hospitalization for PEX

## METHODS


- Retrospective medical chart review at five U.S. pediatric sites
- Data from all children with CF (ages from birth and older) who were **hospitalized** and received **IV antibiotics** for treatment of PEX in 2018 and 2022
- Four elements in REDCap: demographics, admissions, microbiology, inpatient medications

## RESULTS


### Hospitalizations decreased >60% from 2018 to 2022 across our 5 pediatric sites




### No age or sex differences among those hospitalized in 2022 compared with 2018




### Significant racial differences among those hospitalized in 2022 compared with 2018



### Children hospitalized in 2022 were more likely to have two non-F508del mutations



### Changes in lung function were similar in 2018 and 2022



No differences in the proportion of individuals who returned to ≥ 90% or ≥ 100% of baseline ppFEV1

### Other salient study findings

- Lengths of hospitalizations were similar in 2018 and 2022 (mean 11 days vs 10 days,  $p=0.2$ )
- No differences in the percentages of those hospitalized infected with *P. aeruginosa* and methicillin-susceptible *S. aureus* between 2018 and 2022
- Lower percentage of hospitalized children in 2022 (5.5%) were infected with methicillin-resistant *S. aureus* (MRSA) than in 2018 (14%) ( $p=0.018$ )
- Higher percentage of hospitalized children in 2022 (52%) were on a CFTR modulator therapy than in 2018 (34%) ( $p<0.001$ )

## STUDY LIMITATIONS

- Variable practice patterns across sites (diagnosis & treatment)
- Lacking treatment adherence data, particularly around the use of CFTR modulator therapies
- Lacking measures of mental and social health

## CONCLUSIONS

- Reduction in hospitalizations likely reflects the benefit of ETI therapy, as a higher percentage of children hospitalized in 2022 had 2 non-F508del mutations and were not eligible for ETI
- Higher percentage of those hospitalized in 2022 identified as being from minority racial backgrounds, further highlighting the importance of health equity efforts in CF
- Lower percentage of children hospitalized in 2022 were infected with MRSA, concurrent with decreased rates of MRSA community infections in 2022

## ACKNOWLEDGEMENT

2023 CFF Student Traineeship Award (AHMED23H0)