

Low-dose ketamine for acute pain: A Narrative Review



Robert Fuller, ENS Evan Kikla, Andrew Fawcett, LTC Michael April, LTC Michael Patzkowski, Brig Gen (Ret) Kathleen Flarity, COL (Ret) Sean Keenan, COL Vikhyat Bebarta, LTC Steven Schauer

Low-dose ketamine is as effective as opioids for the treatment of acute pain and when used in conjunction can provide a significant opioid sparing effect with limited side effects.

BACKGROUND

- Respiratory compromise and other side effects associated with opioids present obstacles to their employment in combat medicine.
- Low dose ketamine may be an effective alternative for treatment of acute pain while limiting side effects

OBJECTIVES

- Identify recent literature involving LDK analgesia use in prehospital and emergency room settings .
- Identify further research opportunities for the employment of LDK analgesia in prehospital and emergency room settings.

METHODS

Scoping review using PRISMA-ScR guidelines using PubMed Medline. Excluded studies involving sedation, chronic pain, anti-depressant effects or psychiatric conditions.

249

Potential Studies



65

Met inclusion criteria

Inclusion Criteria:

Prospective or retrospective studies evaluating use of LDK for treatment of acute pain in prehospital or ED settings published < 30 Jan 24

RESULTS

Search Results

Prehospital setting	21 Studies
ED setting	44 Studies
RCTs	33

- LDK is increasingly popular in civilian ED and prehospital settings with multiple dosing regimens and routes of administration, including intranasal, under investigation.
- Studies clearly demonstrate that LDK is non-inferior to morphine or fentanyl when used alone.
- LDK may provide an opioid-sparing effect when used as an adjunct.
- LDK is an effective modality for treatment of acute pain in pediatric and adult setting.
- More research is needed to determine the most effective multi-modal combinations incorporating LDK.
- Side effects of LDK are usually mild and transient and typically do not compromise respiratory or hemodynamic status when dosed to <0.5mg/kg.
- Although rare at low doses, the most common side effects of ketamine administration are psychomimetic.
- There are multiple agents that can be co-administered with LDK to mitigate side effects. More research is needed to determine the best options.

CONCLUSIONS

- LDK may offer comparable analgesia to opiates with an advantageous hemodynamic profile when used for moderate-to-severe pain in resource-limited settings.
- Room for further study involving LDK in conjunction with other analgesic options as well as better characterization of dosing regimens for IM and IN routes of administration.

LIMITATIONS

- All studies were limited by substantial heterogeneity in dosing regimen, routes of administration, clinical end points.
- Pre-hospital studies tended to suffer from a lack of RCTs and low population size.
- Limited studies evaluating the effectiveness of LDK in conjunction with specific opioids compared to opioids alone.

DISCLAIMERS

The opinions expressed in this poster presentation are those of the authors and do not reflect the official policy or position of the US Army Medical Department, Department of the Army, Department of Defense, or the US Government.



University of Colorado
Anschutz Medical Campus