

# Impact of Social Vulnerability on Aortic Arch Surgery

Ananya Shah<sup>1</sup>, Adam Carroll<sup>2</sup>, Alejandro Suarez-Pierre<sup>2</sup>, Jessica Rove<sup>3</sup>, Catherine Velopulos<sup>4</sup>, Muhamed Aftab<sup>3</sup>, T. Brett Reece<sup>3</sup>

<sup>1</sup>University of Colorado School of Medicine

<sup>2</sup>Department of Surgery, University of Colorado School of Medicine

<sup>3</sup>Division of Cardiothoracic Surgery, University of Colorado Anschutz Medical Campus

<sup>4</sup>Division of GI, Trauma, and Endocrine Surgery, University of Colorado Anschutz Medical Campus

## Objective:

Socioeconomic status is a known factor influencing morbidity and mortality. The CDC's social vulnerability index (SVI) quantifies neighborhood-level vulnerability. Although high SVI has been associated with adverse outcomes in trauma, CABG, and AAA repair, its impact on aortic arch surgery remains unknown.

## Hypothesis:

In total aortic arch (TAR) and hemiarch (HAR) replacement, higher SVI will correlate with increased comorbidities, surgical acuity, and post-operative morbidity and mortality.

## Methods:

This single-center retrospective study examined adult patients undergoing HAR and TAR performed for aortic aneurysm and/or dissection between 2010 and 2022. SVI was calculated by the patients' residential ZIP codes. The cohort was stratified into three groups: SVI<0.33, SVI 0.33-0.66, and SVI≥0.66. The primary endpoint was postoperative mortality, with control for confounding bias through multivariable logistic regression.

## Results:

For 753 patients, higher SVI correlated with more baseline comorbidities and minority status. SVI was associated with dissection pathology (p=0.04) and urgent/emergent procedures (p=0.02). High SVI was associated with TAR (p=0.01), reflected by lower nadir bladder temperatures (p=0.01), longer cardiopulmonary bypass (p=0.02), longer circulatory arrest times (p=0.01), and more coagulation product usage.

High SVI patients had longer length of stay, higher rates of infection (p=0.01) and a trend towards increased stroke risk. SVI did not correlate with in-hospital or late mortality, confirmed by multivariable regression.

## Conclusion:

In socially vulnerable communities, patients who require aortic arch surgery have more comorbidities and present urgently or emergently with dissection pathology, requiring more extensive arch repair. Post-procedure, they have a higher risk of infection and a trend towards more stroke risk, but not higher rates of other adverse outcomes or increased mortality during or after hospital stay.