

## ANTIBIOTIC PROPHYLAXIS REGIMEN

\*\*\* NO routine dental cleanings/invasive procedures (i.e. skin biopsy, other biopsy's, colonoscopies, etc) 2 MONTHS BEFORE or AFTER YOUR SURGERY.\*\*\*

\*\*\* Any question of a dental abscess or infection must be addressed prior to undergoing surgery!\*\*\*

**It is recommended that antibiotic prophylaxis prior to procedures (including dental appointments for routine cleaning) be taken for at least 2 years after surgery and if well tolerated then preferably for the rest of your life!**

### FOR MOST PROCEDURES ANTIBIOTIC PROPHYLAXIS PRESCRIBED ->

- Amoxicillin 2 grams taken 1 hour prior to procedure/routine dental cleaning
- *If allergic to Penicillin or Amoxicillin ->*
  - take Clindamycin 600 mg orally 1 hour prior to procedure/routine dental cleaning

### FOR GENITOURINARY & GASTROINTESTINAL PROCEDURES ->

- Ciprofloxacin 750 mg orally 1 hour prior to procedure

Hayley E. Ennis, M.D.  
Orthopaedic Surgery, Adult Reconstruction  
University of Colorado – School of Medicine



**UCHealth Broomfield Clinic**  
875 W 136<sup>th</sup> Ave, Broomfield, CO 80023  
Office: 303.544.3820  
Fax: 303.544.3830

**UCHealth Broomfield Hospital**  
11820 Destination Dr, Broomfield, CO 80021

**UCHealth Longmont Clinic**  
1925 W Mountain View Ave, 3<sup>rd</sup> Floor  
Longmont, CO 80501  
Office: 720.718.8288  
Fax: 720.718.0969  
**UCHealth Longs Peak Hospital**  
1750 E Ken Pratt Blvd, Longmont, CO 80504

Official Health Care Partner

