Personal Protective Equipment (PPE)

Perform hand hygiene prior to and after donning ANY mask. It is important to NOT touch the outside of the facemask with your hands. If you touch or adjust your facemask, you must immediately perform hand hygiene.

Eye Protection Contact **Droplet Airborne** Surgical/Procedural Mask **Powered Air Purifying Disposable Gown N95 Respirator Reusable Goggles Respirator (PAPR)** Extended use and reuse: Extended use and reuse: Extended use and reuse: 1 per staff member per shift if clinically indicated (can Use for multiple HCW. Use for multiple HCW. Extended use and reuse: use with multiple patients). Clean and disinfect hood and PAPR · Clean and disinfect after each use. Store in paper bag with employee name and date device after each use. · Discard if damaged. Disposable gowns to be prioritized for aerosolwhen not in use. · Discard hood if torn, visibly soiled or generating procedures, care activities where splashes Place in collection bin at end of shift. and sprays or contact with body fluids are anticipated, damaged. Personal eyeglasses do not provide protection. Discard if contaminated with body fluids or and patients in contact isolation precautions. Extended use and reuse: compromised (damaged/ difficult to breathe through). 1 per staff member per shift (can use with Prioritize for procedures likely to generate Recommended for reuse by multiple HCWs for low- or multiple patients) **Facemasks** respiratory aerosols (bronch, intubation, sputum moderate-risk scenarios and for patients on contact Half Facepiece Reusable Store in paper bag with employee name induction, nebulizer treatment). precautions for MRSA. VRE or MDRO. and date when not in use Respirator Place in reprocessing bin at end of shift Low-risk scenarios: When caring for suspected or confirmed COVID-19 Discard if contaminated with body fluids patients requiring aerosol generating procedures: Basic patient care or cover gown for visitors. or compromised (damaged/difficult to Limit personnel in room. breathe through). Wear N95 for procedure/treatment. Moderate-risk scenarios: DO NOT store and reuse.

- Performing blood draws, suturing, establishing IV access, wound/ostomy care.
- Store in a manner to prevent contamination close to the patient door in the patient room.

Not recommended for reuse if soiled or damaged or if used in high risk scenarios:

High-risk scenarios:

- Precautions for suspected or confirmed COVID-19
- Invasive procedures or surgery
- Lengthy, fluid intense procedures
- Trauma cases
- Special contact precautions
- Precautions for CRE

Discard after use.

When caring for suspected or confirmed **COVID-19 patients NOT requiring aerosol generating** procedures:

- Wear surgical/procedural mask across all patient encounters.
- DO NOT store and reuse.
- Place in bin for reprocessing at each break and at end of day.
- Obtain new mask.

Patient use: (e.g. for transport)

- Provide 1 over the course of hospitalization.
- Store in paper bag in patient room with patient name and date when not in use.
- Discard when no longer needed.



Place in reprocessing bin after procedure/treatment.

urgical mask (Droplet ombo mask/eve shi (N95) + eye shield (Airborne + Eye)

Extended use and reuse:

- Single HCW only.
- Perform negative/positive check prior to use.
- Clean and disinfect device after each use (avoid moistening filters).
- Wipe the plastic where it comes into contact with skin with an alcohol wipe.

Extended use and reuse:

- Use for multiple HCW.
- Clean and disinfect after each use.
- Place in reprocessing bin at end of shift.
- Discard if torn or damaged.

Face Mask w/ Shield is single user only (can use with multiple patients).

